

# EDURANT<sup>®</sup> Tablets

*rilpivirine*

---

## Consumer Medicine Information

---

### What is in this leaflet

This leaflet answers some common questions about EDURANT tablets. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you being given EDURANT against the benefits this medicine is expected to have for you.

**If you have any concerns about being given EDURANT ask your doctor.**

**Keep this leaflet while you are taking EDURANT.** You may need to read it again.

---

### What EDURANT is used for

EDURANT is a medicine used for the treatment of Human Immunodeficiency Virus (HIV) infection. It belongs to a group of anti-HIV medicines called non-nucleoside reverse transcriptase inhibitors (NNRTIs). EDURANT works by reducing the amount of HIV in your body. This will improve your immune system and reduce the risk of developing illnesses linked to HIV infection.

EDURANT in combination with other HIV medicines is used to treat adults who are infected by HIV and who have not used other anti-HIV medicines before.

EDURANT must be taken in combination with other anti-HIV medicines. Your doctor will discuss with you which combination of medicines is best for you.

**Ask your doctor if you have any questions about why EDURANT has been prescribed for you.**

**This medicine is available only with a doctor's prescription.**

---

### Before you take EDURANT

**When you must not use it:**

**Do not take EDURANT:**

- if you are **allergic** (hypersensitive) to rilpivirine, or other ingredients of EDURANT. See **Product Description** at the end of this leaflet for a list of ingredients.

**Do not take EDURANT:**

- if the packaging is torn or shows signs of tampering.
- if the expiry date (month and year) printed on the pack has passed. If you take EDURANT after the expiry date it may not work.

**Do not combine EDURANT with any of the following medicines:**

- carbamazepine, oxcarbazepine, phenobarbital, phenytoin (to treat epilepsy or prevent seizures)
- non-nucleoside reverse transcriptase inhibitors (NNRTIs) (HIV medicines)

- rifampicin, rifapentine (to treat some infections such as tuberculosis)
- products that contain St John's wort (*Hypericum perforatum*) (a herbal product used for depression)
- omeprazole, esomeprazole, lansoprazole, rabeprazole, pantoprazole (to treat gastrointestinal ulcers, heartburn or acid reflux disease)
- dexamethasone (a corticosteroid used in a variety of conditions such as inflammation and allergic reactions)

If you are taking any of these, ask your doctor for advice.

There are other medicines that you need to be careful of when taking EDURANT (see Taking other medicines).

**Before you start to use it:**

**Take special care with EDURANT:**

EDURANT is not a cure for HIV infection. It is part of a treatment reducing the amount of virus in the blood.

People taking EDURANT must continue to use appropriate precautions (a condom or other barrier method) to lower the chance of sexual contact with any body fluids such as semen, vaginal secretions or blood.

People taking EDURANT may still develop infections or other illnesses associated with HIV infection. You must keep in regular contact with your doctor to monitor your health while taking EDURANT.

EDURANT is not for use in paediatric patients, because it has not been sufficiently studied in this patient population.

EDURANT has only been used in a limited number of patients aged 55 years or older. If you belong to this age group, discuss the use of EDURANT with your doctor.

**Tell your doctor if you have or have had any medical conditions, especially the following:**

- **Problems with your liver.** Including hepatitis B and/or C. Your doctor may need to evaluate how severe your liver disease is before deciding if you can take EDURANT.
- **Symptoms of infection or inflammation.** In some patients with advanced HIV infection, signs and symptoms of infection or inflammation may occur after anti-HIV treatment is started. It is believed that these symptoms are due to an improvement in the body's immune response.
- 

**Tell your doctor immediately if you are pregnant or breast-feeding, or intend to become pregnant or breast-feed.**

HIV infected mothers must not breast-feed, as there is a possibility of infecting the baby with HIV.

During pregnancy and breast-feeding you should not take EDURANT, unless it is specifically approved by your doctor.

**If you have not told your doctor about any of the above, tell them before you start treatment with EDURANT.**

## **Taking other medicines:**

**EDURANT might interact with other medicines. Tell your doctor if you are taking any other medicines, including medicines you can buy without a prescription from a pharmacy, supermarket or health food shop.**

EDURANT can be combined with most other anti-HIV medicines belonging to another class. However, some combinations are not recommended. In other cases, increased monitoring and/or a change in the dose of the medicine may be needed. Therefore, always tell your doctor which other anti-HIV medicines you take. Carefully follow your doctor's instructions, on which medicines can be combined.

In addition to other anti-HIV medicines that you take, tell your doctor or pharmacist if you are taking any of the following:

- rifabutin (a medicine to treat some bacterial infections). If you take this medicine while taking EDURANT, please carefully read the section entitled 'Taking EDURANT'
- cimetidine, famotidine, nizatidine, ranitidine (antihistamines used to treat stomach or intestinal ulcers or used to relieve heartburn from acid reflux)
- antacids (for example, aluminium / magnesium hydroxide, calcium carbonate)
- clarithromycin, erythromycin, troleandomycin (antibiotics)
- methadone (strong pain killer also used in drug dependence programs)

---

## **Taking EDURANT**

---

### **How much EDURANT to take:**

Always take EDURANT exactly as your doctor has told you. You must check with your doctor if you are not sure.

**The usual dose of EDURANT is one tablet taken once daily with a meal.**

### **Instructions:**

- It is important that you take EDURANT with a meal. A meal is important to get the right drug levels in your body. A protein drink alone does not replace a meal.

If you take rifabutin (a medicine to treat some bacterial infections), take two tablets of EDURANT once a day. When you stop taking rifabutin, take one tablet of EDURANT once a day. Talk to your doctor or pharmacist if you are not sure.

If you take an antacid (a medicine to treat diseases related to the acid in the stomach such as aluminium/magnesium hydroxide, calcium carbonate), take the antacid either at least 2 hours before or at least 4 hours after EDURANT. If you take an H<sub>2</sub>-receptor antagonist (medicines used for stomach ulcers, heartburn, or acid reflux disease (such as cimetidine, famotidine, nizatidine or ranitidine), take the H<sub>2</sub>-receptor antagonist at least 12 hours before or at least 4 hours after EDURANT. Importantly, proton pump inhibitors (such as omeprazole, lansoprazole, rabeprazole, pantoprazole, esomeprazole) also available for these conditions should not be taken with EDURANT.

Make sure that you always have enough EDURANT available so

that you don't run out. For example, in case you cannot return home, need to travel or stay in a hospital.

### Removing the child resistant cap



The plastic bottle comes with a child resistant cap and should be opened as follows:

- Push the plastic screw cap down while turning it counter-clockwise.
- Remove the unscrewed cap.

### What do I do if I forget to take EDURANT?

If you notice within **12 hours** of the time you usually take EDURANT, you must take the tablet as soon as possible. Always take the tablet with a meal. Then take the next dose as usual. If you notice after **12 hours**, then skip the intake and take the next doses as usual. Do not take a double dose to make up for a forgotten dose.

### What do I do if I take too much? (overdose):

**If you think you or anybody else has taken too much EDURANT, contact your doctor, pharmacist or the Poisons Information Centre who will advise you what to do.**

You can contact the Poisons Information Centre by dialling:

- Australia: 13 11 26
- New Zealand: 0800 POISON or 0800 764 766.

---

## While you are taking EDURANT

---

### Things you must do:

#### **Do not stop taking EDURANT without talking to your doctor first**

HIV therapy may increase your sense of well being. Even if you feel better, do not stop taking EDURANT or your other anti-HIV medicines. Doing so could increase the risk of the virus developing resistance. Talk to your doctor first.

#### **Be sure to keep all your doctor's appointments so your progress can be checked.**

Your doctor will want to do some blood and other tests from time to time to check on your progress.

#### **Be sure to follow up your doctor's instructions about other medicines you should take, and other things you should do.**

Ask your doctor or pharmacist if you have any questions.

#### **Tell any other doctors and pharmacists who are treating you that you are taking EDURANT. If you are undergoing anaesthesia, tell your anaesthetist that you are taking EDURANT.**

**If you are about to be started on any new medicines, tell your doctor or pharmacist that you are taking EDURANT.**

**If you become pregnant while taking EDURANT, tell your doctor immediately.**

If you have any further questions on the use of this product, ask your doctor.

### Things to be careful of

#### **Driving and using machines**

EDURANT has no or negligible influence on the ability to drive and use machines.

---

## Side Effects

---

Like all medicines, EDURANT can have side effects. Some of these effects may be serious.

#### **Tell your doctor or pharmacist if you do not feel well while you are being treated with EDURANT.**

When treating HIV infection, it is not always easy to identify what side effects are caused by EDURANT, which are caused by other medicines you are taking, or which are caused by the HIV infection itself.

The most common side effects are:

- skin rash. The rash is usually mild to moderate. In rare instances, rash can be potentially life-threatening. It is therefore important to contact your doctor immediately if you develop a rash. Your doctor will advise you how to deal with your symptoms or whether EDURANT must be stopped.
- stomach pain, nausea
- depression, difficulty falling asleep (insomnia), abnormal dreams, sleep disorders
- decreased appetite
- tiredness
- changes in your routine liver tests
- headache

Uncommon side effects include:

- changes in your routine blood tests
- dizziness, drowsiness
- vomiting, stomach discomfort

Some side effects are typical for combination anti-HIV therapy. These are:

- change in body shape due to fat redistribution. This may include loss of fat from legs, arms and face, increased fat in the abdomen (belly) and other internal organs, breast enlargement and fatty lumps on the back of the neck (buffalo hump). The cause and long-term health effects of these conditions are not known at this time.
- immune reconstitution syndrome. In some patients with advanced HIV infection (AIDS) and a history of opportunistic infection, signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started, including EDURANT. In addition to the opportunistic infections, autoimmune disorders (a condition that occurs when the immune system attacks healthy body tissue) may also occur after you start taking medicines for the treatment of your HIV infection. Autoimmune disorders may occur many months after the start of treatment. It is believed that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms.

If you experience any of these side effects and they worry you, or if you notice any side effects not listed in this leaflet, please tell your doctor.

Other side effects not listed above may also occur in some people.

---

## Product Description

---

### Storage

EDURANT tablets should be kept out of reach of children, in a location where the temperature stays below 30°C.

EDURANT tablets should be stored in the original bottle. Keep the bottle tightly closed in order to protect from moisture and light.

### What it looks like:

EDURANT tablets are white to off-white, film-coated, round, biconvex tablet, with "TMC" on one side and "25" on the other side. Each plastic bottle contains 30 tablets.

### Ingredients

Active ingredient:

- rilpivirine 25 mg

Other ingredients:

- triacetin
- silicified microcrystalline cellulose
- magnesium stearate
- hypromellose
- croscarmellose sodium
- lactose monohydrate

- povidone
- polysorbate 20
- titanium dioxide
- macrogol 3000

### Sponsor

JANSSEN-CILAG Pty Ltd  
1-5 Khartoum Rd  
Macquarie Park NSW 2113  
Australia  
Telephone: 1800 226 334

NZ Office: Auckland, New Zealand

Telephone: 0800 800 806

**Tibotec, a division of  
JANSSEN-CILAG Pty  
Ltd**



### Registration number

25 mg Tablet: AUST R 176666

This leaflet was prepared in May 2024.

® EDURANT is a registered trademark of Tibotec Pharmaceuticals Ltd.