

Oxycodone Hydrochloride Tablets, USP (Immediate Release)

Oxycodone hydrochloride (ox-ee-code-own hi-dro-klor-ide) (USP)

Immediate release tablets 5mg, 10mg and 20mg

What is in this leaflet

Please read this leaflet carefully before you start using oxycodone hydrochloride immediate release tablets.

This leaflet answers some common questions about oxycodone hydrochloride immediate release tablets. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking this medicine against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine. You may need to read it again.

What oxycodone hydrochloride immediate release tablets are used for

Oxycodone hydrochloride immediate release tablets contain oxycodone hydrochloride. Oxycodone belongs to a group of medicines called opioid analgesics.

oxycodone hydrochloride immediate release tablets are used to relieve moderate to severe pain.

Your doctor, however, may have prescribed it for another reason.

Ask your doctor if you have any questions about why it has been prescribed for you.

As with all strong painkillers, your body may become used to you taking oxycodone hydrochloride immediate release tablets. Taking it may result in physical dependence. Physical dependence means that you may experience withdrawal symptoms if you stop taking oxycodone suddenly, so it is important to take it exactly as directed by your doctor.

This medicine is only available with a doctor's prescription.

Before you take oxycodone hydrochloride immediate release tablets

Long-term use of oxycodone hydrochloride immediate release tablets may result in a decrease in sex hormone levels which may affect sperm production in men and the menstrual cycle in females. Talk to your doctor if you have any concerns.

When you must not take it

Do not take oxycodone hydrochloride immediate release tablets if you:

- suffer from shallow breathing or have any breathing problems such as acute asthma, chronic bronchitis or impaired lung function
- are severely drowsy or have a reduced level of consciousness
- suffer from irregular or fast heartbeats or changes in the way the heart beats
- have heart problems due to long-term lung disease
- have just consumed a large amount of alcohol, regularly consume large amounts of alcohol or have confusion and shaking due to alcohol withdrawal
- suffer from convulsions, fits or seizures
- have a head injury, brain tumour or have raised pressure within the head, brain or spinal cord
- have sudden severe abdominal pain
- have a condition where your stomach empties more slowly than it should, or your small bowel does not work properly
- are about to have an operation (including surgery on your spine for pain relief in the next 6 hours)
- have severe kidney or liver disease
- take a medicine for depression called a 'monoamine oxidase inhibitor' or have taken any in the last two weeks.

Do not take oxycodone hydrochloride immediate release tablets if you are allergic to oxycodone, opioid painkillers, or any of the ingredients listed at the end of this leaflet.

Do not take this medicine after the expiry date (EXP) printed on the pack.

If you take it after the expiry date has passed, it may not work very well.

Do not take it if the packaging shows signs of tampering.

Do not take this medicine if you are pregnant or intend to become pregnant whilst taking this medicine.

Like most medicines of this kind, oxycodone hydrochloride immediate release tablets are not recommended to be taken during pregnancy. Your doctor will discuss the risks and benefits of taking them if you are pregnant.

Do not give this medicine to a child younger than 18 years of age.

Safety and effectiveness in children younger than 18 years of age have not been established.

If you are not sure whether you should start taking oxycodone hydrochloride immediate release tablets, talk to your doctor.

Before you start to take it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have had any medical conditions, especially the following:

- low blood pressure
- increased prostate size or difficulty passing urine
- chronic lung, liver or kidney disease
- disease of your gall bladder or bile duct
- inflammation of the pancreas
- underactive adrenal glands
- underactive thyroid gland
- inflammatory bowel disease or recent abdominal surgery
- severe mental condition involving losing contact with reality, hearing voices or an inability to think clearly
- an addiction or history of abuse of alcohol, opioids or other drugs.

This medicine is not recommended to be taken during labour.

Oxycodone given to the mother during labour can cause breathing problems and signs of withdrawal in the newborn.

Tell your doctor if you are breastfeeding or planning to breastfeed.

Oxycodone can pass into the breast milk and can harm the baby. Your doctor can discuss with you the risks and benefits involved.

If you have not told your doctor about any of the above, tell them before you take any oxycodone hydrochloride immediate release tablets.

Taking other medicines

Tell your doctor if you are taking any other medicines or dietary supplements, including any that you buy without a prescription from your pharmacy, supermarket or health food shop. You should also tell any health professional who is prescribing a new medication for you that you are taking oxycodone hydrochloride immediate release tablets.

Some medicines, dietary supplements and oxycodone hydrochloride immediate release tablets may interfere with each other. These include:

- medicines to treat depression, psychiatric or mental disorders
- medicines to treat depression belonging to a group called monoamine oxidase inhibitors must be stopped 14 days before oxycodone hydrochloride immediate release tablets are taken
- medicines to help you sleep
- medicines to put you to sleep during an operation or procedure
- medicines to relax your muscles

- medicines to lower blood pressure
- quinidine and other medicines to treat the heart
- medicines to treat convulsions, e.g. phenytoin and carbamazepine
- medicines to thin the blood e.g. coumarin derivatives such as warfarin
- cimetidine, a medicine to treat stomach ulcers and heartburn
- medicines to relieve stomach cramps or spasms, or to prevent travel sickness and the symptoms of allergies
- medicines to treat Parkinson's disease
- medicines to treat urinary incontinence
- medicines to stop nausea or vomiting e.g. metoclopramide
- other pain relievers including other opioids
- medicines to treat fungal infections, e.g. ketoconazole
- antibiotics, e.g. clarithromycin, rifampin
- medicines to treat HIV infection and AIDS, e.g. ritonavir
- alcohol
- St John's wort (a herbal preparation)
- grapefruit and grapefruit juice.

These medicines, dietary supplements or alcohol may be affected by oxycodone hydrochloride immediate release tablets or may affect how well oxycodone hydrochloride immediate release tablets work or may increase side effects. You may need to use different amounts of your medicines, or take different medicines.

Your doctor or pharmacist has more information on medicines and dietary supplements to be careful with or avoid while taking this medicine.

How to take oxycodone hydrochloride immediate release tablets

How much to take

Your doctor will tell you exactly how much to take.

Follow the instructions given to you by your doctor or pharmacist exactly.

How to take it

Swallow oxycodone hydrochloride immediate release tablets with a full glass of water or other fluid.

You must only take oxycodone hydrochloride immediate release tablets by mouth.

Taking this medicine in a manner other than that prescribed by your doctor can be harmful to your health.

When to take it

Take oxycodone hydrochloride immediate release tablets every 4 to 6 hours or as directed by your doctor.

If you begin to experience pain and you are taking your oxycodone hydrochloride immediate release tablets as prescribed, contact your doctor as your dosage may have to be reviewed.

How long to take it

Continue taking your medicine for as long as your doctor tells you.

If you stop taking this medicine suddenly, your pain may worsen and you may experience withdrawal symptoms such as:

- body aches
- loss of appetite, nausea, stomach cramps or diarrhoea
- fast heart rate
- sneezing or runny nose
- chills, tremors, shivering or fever
- trouble sleeping
- increased sweating and yawning
- weakness
- nervousness or restlessness.

If you forget to take it

If you forget to take your tablets, contact your doctor or pharmacist for advice.

Do not take a double dose to make up for the dose you have missed.

This will increase the chance of you getting unwanted side effects.

While you are taking oxycodone hydrochloride immediate release tablets

Things you must do

Take oxycodone hydrochloride immediate release tablets exactly as your doctor has prescribed.

Before you start on a new medicine, remind your doctor and pharmacist that you are taking oxycodone hydrochloride immediate release tablets.

Tell any other doctors, dentists and pharmacists who treat you that you are taking this medicine.

If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine.

It may affect other medicines used during surgery.

If you become pregnant while taking this medicine, tell your doctor immediately.

Keep all of your doctor's appointments so that your progress can be checked.

Tell your doctor if your pain is getting worse. Always discuss any problems or difficulties you have while you are taking oxycodone hydrochloride immediate release tablets.

Tolerance to oxycodone may develop which means that the effect of the medicine may decrease. If this happens, your doctor may review your dose so that you get adequate pain relief.

Keep enough oxycodone hydrochloride immediate release tablets with you to last over weekends and holidays.

Things you must not do

Do not drink alcohol while you are taking oxycodone hydrochloride immediate release tablets.

Drinking alcohol whilst taking oxycodone hydrochloride immediate release tablets may make you feel more sleepy and increase the risk of serious side effects such as shallow breathing with the risk of stopping breathing and loss of consciousness.

Do not take oxycodone hydrochloride immediate release tablets to treat any other complaint unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not stop taking your medicine, exceed the dose recommended or change the dosage without checking with your doctor.

Over time your body may become used to oxycodone. If you stop taking it suddenly, your pain may worsen and you may have unwanted side effects such as withdrawal symptoms. This is called physical dependence.

If you need to stop taking this medicine, your doctor will gradually reduce the amount you take each day, if possible, before stopping the medicine completely.

Things to be careful of

Do not drive or operate machinery until you know how oxycodone hydrochloride immediate release tablets affects you.

Oxycodone hydrochloride immediate release tablets may cause drowsiness, dizziness, hallucinations, disorientation, blurred vision or other vision problems or may affect alertness. If you are affected you should not drive or operate machinery. Discuss these effects with your doctor.

Be careful if you are elderly, unwell or taking other medicines.

Some people may experience side effects such as drowsiness, confusion, dizziness and unsteadiness, which may increase the risk of a fall.

If you feel light-headed, dizzy or faint when getting out of bed or standing up, get up slowly.

Standing up slowly will help your body get used to the change in position and blood pressure. If this problem continues or gets worse, talk to your doctor.

Tell your doctor if you suffer from nausea or vomiting when taking oxycodone hydrochloride immediate release tablets.

If you vomit after your dose, your pain may come back, as you may not have absorbed your medicine. If this happens, speak to your doctor. Your doctor may prescribe some medicine to help you stop vomiting.

Tell your doctor if taking oxycodone hydrochloride immediate release tablets causes constipation.

Your doctor can advise about your diet, the proper use of laxatives and suitable exercise you can do to help manage this.

There is potential for abuse of oxycodone and the development of dependence. It is important that you discuss this issue with your doctor.

In case of overdose

If you take too much (overdose)

Immediately telephone your doctor or the National Poisons Centre (telephone 0800 POISON or 0800 764 766), or go to accident and emergency at your nearest hospital, if you think that you or anyone else may have taken too many oxycodone hydrochloride immediate release tablets.

Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

Keep telephone numbers for these places handy.

You should also follow the above steps if someone other than you has accidentally taken the tablets that were prescribed for you.

If someone takes an overdose they may experience difficulties in breathing, become drowsy and tired, lack muscle tone, have cold or clammy skin, have constricted pupils, have very low blood pressure or slow heart rate, and possibly may even become unconscious or die.

When seeking medical attention, take this leaflet and any remaining tablets with you to show the doctor. Also tell them about any other medicines or alcohol which have been taken.

Side effects

All medicines may have some unwanted side effects. Sometimes they are serious but most of the time they are not. You may need medical treatment if you get some of the side effects.

Your doctor has weighed the risks of this medicine against the benefits they expect it will have for you.

Do not be alarmed by this list of possible side effects.

Not everybody experiences them.

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking oxycodone hydrochloride immediate release tablets.

This medicine helps people with moderate to severe pain, but it may have unwanted side effects in some people. Other side effects not listed here may also occur in some people.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if you notice any of the following and they worry you:

- mild abdominal symptoms such as diarrhoea, feeling sick (nausea), decreased appetite or constipation
- dry mouth, hiccups, sore throat, trouble swallowing or changes in voice
- excessive sweating
- feeling anxious or nervous, trouble sleeping or abnormal dreams
- trouble with your balance, vertigo
- new problems with your eyesight
- skin rash, itching, chills or fever
- muscle problems, such as spasms, twitching or tremors
- swelling of legs or ankles
- absence of menstrual periods
- impotence
- decreased sexual drive.

Tell your doctor as soon as possible if you notice any of the following and they worry you:

- stomach discomfort, vomiting, indigestion or abdominal pain
- abnormal thinking, changes in mood or feeling deep sadness
- drowsiness, fainting or dizziness especially when standing up
- slow or noticeable heartbeats
- headache or confusion
- unusual weakness or loss of strength or trouble walking
- fatigue, feeling of tiredness, drowsiness or lack of energy
- changes in passing urine such as the volume passed, pain or
- feeling the need to urinate urgently.
- increased levels of pain

The above list includes serious side effects that may require medical treatment.

If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- your breathing slows or weakens
- you have an allergic reaction: shortness of breath, wheezing, shallow or difficult breathing; swelling of the face, lips, tongue, throat or other parts of the body; rash, itching or hives on the skin
- seizures, fits or convulsions
- fast or irregular heartbeats
- chest pain or chest tightness

- temporarily stopping breathing while you sleep.

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

When seeking medical attention, take this leaflet and any remaining tablets with you to show the doctor.

After taking oxycodone hydrochloride immediate release tablets

Storage

Keep your tablets in the bottle until it is time to take them.

If you take the tablets out of the bottle they may not keep as well.

Keep your tablets in a cool, dry place, where the temperature stays below 25°C.

Do not store it or any other medicine in the bathroom, near a sink or on a window sill.

Do not leave it in the car on hot days

Heat and damp can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking the tablets or the tablets have passed their expiry date, ask your pharmacist how to dispose of medicines no longer required.

Product description

What it looks like

Oxycodone hydrochloride immediate release tablets come in three strengths:

5 mg tablets: white, round, biconvex tablets debossed with “A” on the left and “04” on the right of the score on one side and plain on the other side.

10 mg tablets: pink, round, biconvex tablets debossed with “A” on the left and “48” on the right of the score on one side and plain on the other side.

20 mg tablets: grey, round, biconvex tablets debossed with “A” on the left and “50” on the right of the score on one side and plain on the other side.

Oxycodone hydrochloride immediate release tablets come in bottles containing 100 tablets.

Ingredients

Active ingredient:

- 5 mg tablets contain 5 mg oxycodone hydrochloride
- 10 mg tablets contain 10 mg oxycodone hydrochloride
- 20 mg tablets contain 20 mg oxycodone hydrochloride

Inactive ingredients:

- corn starch
- lactose monohydrate
- microcrystalline cellulose
- sodium starch glycolate
- stearic acid

The 10 mg tablets also contain the colourant D&C Red No. 27.

The 20 mg tablets also contain the colourants FD&C Blue No. 2, FD&C Red No. 40 and FD&C Yellow No. 6.

This medicine contains lactose.

Sponsor Details

Oxycodone hydrochloride immediate release tablets are supplied in New Zealand by:

Wellmed.NZ Limited
Level 1
50 Customhouse Quay
Wellington 6011
NEW ZEALAND
Phone: 021 048 8867

Date of Preparation

This leaflet was prepared on 12 July 2024.