

## VANCOMYCIN VIATRIS

### *Vancomycin Viatris Powder for Infusion 500mg*



---

## What is in this leaflet

---

This leaflet answers some common questions about Vancomycin Viatris.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you being given Vancomycin Viatris against the benefits they expect it will have for you.

**If you have any concerns about being given this medicine, ask your doctor or pharmacist.**

**Keep this leaflet.** You may need to read it again.

---

## What Vancomycin Viatris is used for

---

This medicine is used to treat severe infections caused by bacteria, such as:

- staphylococcal infections (including methicillin resistant staphylococci)
- endocarditis (an infection of the lining of the heart)
- osteomyelitis (bone infection)
- pneumonia (lung infection)
- septicaemia (infection of the blood)
- infections of the skin or the skin structure
- enterocolitis (infection in the intestines).

This medicine belongs to a group of medicines called glycopeptide antibiotics.

**Ask your doctor if you have any questions about why this medicine has been prescribed for you.**

Your doctor may have prescribed it for another reason.

This medicine is not addictive.

This medicine is available only with a doctor's prescription.

---

## Before you are given Vancomycin Viatris

---

### *When you must not be given it*

**You must not be given Vancomycin Viatris if you have an allergy to:**

- any medicine containing Vancomycin
- any of the ingredients listed at the end of this leaflet
- other antibiotics that are the same type.

Some of the symptoms of an allergic reaction may include: shortness of breath; wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin.

**If you are not sure whether you should be given this medicine, talk to your doctor.**

### *Before you are given it*

**Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.**

**Tell your doctor if you have or have had any of the following medical conditions:**

- kidney problems
- hearing problems

- inflammatory bowel disorders.

**Tell your doctor if you are pregnant or plan to become pregnant or are breastfeeding.**

Your doctor can discuss with you the risks and benefits involved.

**If you have not told your doctor about any of the above, tell him/her before you are given Vancomycin Viatris.**

### *Taking other medicines*

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines and Vancomycin Viatris may interfere with each other. These include:

- some other medicines used to treat infections, such as amikacin, amphotericin B, streptomycin, kanamycin, colistin, gentamicin, neomycin, tobramycin and piperacillin/tazobactam
- cisplatin, an anti-cancer medicine
- some fluid tablets (diuretics), such as ethacrynic acid and frusemide
- anaesthetic agents
- muscle relaxants
- cholestyramine, a powder taken to lower cholesterol levels
- suxamethonium or vecuronium, medicines used to relax muscles.

These medicines may be affected by Vancomycin Viatris or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to

be careful with or avoid while taking this medicine.

---

## How Vancomycin Viatris is given

---

### *How much is given*

Your doctor will decide what dose of Vancomycin Viatris you will receive and how long you will receive it for. This depends on your condition and other factors, such as your weight and kidney function.

### *How it is given*

Vancomycin Viatris is usually given as a slow injection into a vein (intravenous drip). Vancomycin Viatris (as hydrochloride), powder for Intravenous Infusion must only be given by a doctor or nurse.

Sometimes, Vancomycin Viatris is taken orally (swallowed). Vancomycin Viatris may be diluted with water and flavoured with syrup (as it has an unpleasant taste) and given as a drink.

---

## While you are being treated with Vancomycin Viatris

---

### *Things you must do*

**If you are about to be started on any new medicine, remind your doctor and pharmacist that you are being given Vancomycin Viatris.**

**Tell any other doctors, dentists, and pharmacists who treat you that you are being given this medicine.**

**If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine. It may affect other medicines used during surgery.**

**Keep all of your doctor's appointments so that your progress can be checked.** Your doctor may do some tests such as leukocyte count, and

hearing tests from time to time to make sure the medicine is working and to prevent unwanted side effects.

**If you become pregnant soon after being given Vancomycin Viatris, tell your doctor immediately.**

### *Things to be careful of*

**Be careful driving or operating machinery until you know how Vancomycin Viatris affects you.**

This medicine may cause dizziness in some people. If this occurs, do not drive, operate machinery or do anything else that could be dangerous. If you drink alcohol, dizziness may be worse.

---

## In case of overdose

---

As Vancomycin Viatris is usually given to you in hospital under the supervision of your doctor, it is very unlikely that you will receive an overdose.

**However, if you experience severe side effects after being given Vancomycin Viatris, tell your doctor or nurse immediately.**

Symptoms of a Vancomycin Viatris overdose may include the side effects listed below in the 'Side Effects' section, but are usually of a more severe nature.

**Immediately contact your doctor or the National Poisons Information Centre (0800 POISON or 0800 764 766) or go to the Emergency department at your nearest hospital, if you think that you may have been given or had too much, or if anyone else has taken any Vancomycin Viatris by mistake. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.** Take the container of Vancomycin Viatris with you if you can.

**Ask your doctor or pharmacist if you have any concerns.**

---

## Side effects

---

**Tell your doctor or nurse as soon as possible if you do not feel well while you are being treated with Vancomycin Viatris.**

This medicine helps most people with certain serious infections, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

If you are over 65 years of age you may have an increased chance of getting side effects.

**Do not be alarmed by the following lists of side effects. You may not experience any of them.**

**Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor or nurse if you notice any of the following and they worry you:**

- irritation at the injection site
- nausea, vomiting or mild diarrhoea
- dizziness.

The above list includes side effects that are usually mild and short-lived.

**Tell your doctor or nurse immediately if you notice any of the following:**

- signs of an allergic reaction, such as shortness of breath, wheezing or difficulty breathing, swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin
- rare skin condition with severe blisters and bleeding in the lips, eyes, mouth, nose and genitals
- swelling and redness along a vein which is extremely tender when touched
- chills or fever, which may be accompanied by shivering
- flushing of the upper body or pain and muscle spasm of the chest and back

- fast, slow or irregular heart beat, heart palpitations
- ringing in the ears or hearing loss, dizziness or problems with your balance
- other infections
- bruising.

The above list includes serious side effects that may require medical attention. Serious side effects are rare.

## ***After finishing it***

**Tell your doctor immediately if you notice any of the following side effects, particularly if they occur several weeks after stopping treatment with Vancomycin Viatris:**

- severe abdominal cramps or stomach cramps
- watery and severe diarrhoea which may also be bloody
- fever, in combination with one or both of the above.

These are serious side effects. You may have a serious condition affecting your bowel. Therefore, you may need urgent medical attention. However, these side effects are rare.

**Tell your doctor or nurse if you notice anything that is making you feel unwell.** Other side effects not listed above may occur in some patients. Some of these side effects (for example changes in kidney function) can only be found when your doctor does tests from time to time to check your progress.

---

## **After using Vancomycin Viatris**

---

### ***Storage***

Vancomycin Viatris will be stored in the pharmacy or on the ward. The injection is kept in a cool dry place, protected from light, where the temperature stays below 25°C.

---

## **Product description**

---

### ***What it looks like***

Vancomycin Viatris Powder for Infusion is an off white to light tan coloured lyophilized plug or powder for reconstitution, which has been prepared in a sterile fashion. When reconstituted in water, it is a clear solution with a pH of 2.5 – 4.5.

### ***Ingredients***

#### Active ingredient:

Vancomycin Viatris 500 mg contains 500 mg of Vancomycin Viatris hydrochloride as the active ingredient.

#### Inactive ingredients:

Vancomycin Viatris also contains:

- hydrochloric acid
- sodium hydroxide

---

## **If you want to know more**

---

Should you have any questions regarding this product, please contact your pharmacist or doctor.

### ***Who supplies this medicine***

Vancomycin Viatris is supplied in New Zealand by:

Viatri Ltd  
PO Box 11-183  
Ellerslie  
AUCKLAND  
[www.viatris.co.nz](http://www.viatris.co.nz)  
Telephone 0800 168 169

---

## **Date of Preparation**

---

15 August 2025 (Based on datasheet dated 15 August 2025).