

## ENTACAPONE VIATRIS

### *Entacapone Film-coated Tablets 200 mg*



---

## What is in this leaflet

---

Please read this leaflet carefully before you start taking Entacapone Viatrix.

This leaflet answers some common questions about Entacapone Viatrix.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Entacapone Viatrix against the benefits they expect it will have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.** You may need to read it again.

---

## What Entacapone Viatrix is used for

---

Entacapone Viatrix is used to treat the symptoms of Parkinson's disease in people who are already taking levodopa. Levodopa is usually supplied as a combination medicine with carbidopa or benserazide. When levodopa/carbidopa or levodopa/benserazide do not control the symptoms, Entacapone Viatrix can be added.

Parkinson's disease is a disorder of the nervous system. It is caused by a lack of dopamine, a natural substance that is produced in the brain. Dopamine relays messages in the part of the brain that controls muscle movement. When too little dopamine is produced, problems with movement result.

Entacapone Viatrix contains the active ingredient entacapone. This medicine belongs to a group of medicines called catechol-O-methyl transferase (COMT) inhibitors.

Entacapone Viatrix is always used together with levodopa. Levodopa works by increasing the level of dopamine in the brain and Entacapone Viatrix works by making the effect of levodopa last longer. Together, Entacapone Viatrix and levodopa help to relieve symptoms such as shaking of the limbs, stiffness and slowness of movement, which make it difficult to perform normal daily activities. Other medicines can also be added to help treat this condition.

**Ask your doctor if you have any questions about why this medicine has been prescribed for you.**

Your doctor may have prescribed it for another reason.

This medicine is available only with a doctor's prescription.

There is not enough information to recommend the use of this medicine for children under the age of 18 years.

---

## Before you take Entacapone Viatrix

---

**When you must not take it:**

**Do not take Entacapone Viatrix if you have an allergy to:**

- any medicine containing entacapone
- any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include: shortness of breath; wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin.

**Do not take this medicine if you have:**

- liver problems
- phaeochromocytoma, a rare tumour of the adrenal gland which can cause your blood pressure to rise to a dangerous level
- a history of neuroleptic malignant syndrome (NMS), a serious condition with symptoms such as sudden increase in body temperature, sweating, fast heartbeat, muscle stiffness and fluctuating blood pressure
- rhabdomyolysis, a condition with severe muscle weakness that was not due to injury.

If you are not sure whether any of the above conditions apply to you, ask your doctor.

**Do not take Entacapone Viatrix if you are pregnant.**

There is not enough information to recommend its use during pregnancy.

**Do not breastfeed while you are taking this medicine.**

It is not known if the active ingredient in Entacapone Viatrix passes into breast milk and could affect your baby.

**Do not take this medicine after the expiry date printed on the pack or if the packaging is damaged or shows signs of tampering.**

If it has expired or is damaged, return it to your pharmacist for disposal.

**If you are not sure whether you should start taking this medicine, talk to your doctor.**

***Before you start to take it:***

**Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.**

**Tell your doctor if you have or have had any of the following medical conditions:**

- severe kidney disease requiring dialysis
- heart attack or any other heart disease.

**If you have not told your doctor about any of the above, tell him/her before you start taking Entacapone Viatris.**

### ***Taking other medicines***

**Do not take Entacapone Viatris if you are taking a medicine called a monoamine-oxidase inhibitor (MAOI) without first checking with your doctor.**

MAOI medicines are usually used to treat depression. Taking Entacapone Viatris together with some, but not all, MAOI medicines may cause serious side effects. Your doctor will know whether or not the MAOI medicine you are taking can be safely taken with Entacapone Viatris.

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines and Entacapone Viatris may interfere with each other. These include:

- other medicines used for Parkinson's disease
- some medicines used to treat depression
- methyldopa, a medicine for high blood pressure
- warfarin, a medicine used to prevent blood clots
- iron supplements.

These medicines may be affected by Entacapone Viatris or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to

be careful with or avoid while taking this medicine.

---

## **How to take Entacapone Viatris**

---

**Follow all directions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the label, ask your doctor or pharmacist for help.**

### ***How much to take***

The usual dose is one Entacapone Viatris tablet (200 mg) each time you take a dose of levodopa.

People with moderate to severe Parkinson's disease usually take from 4 to 7 tablets of Entacapone Viatris each day.

The maximum dose is 10 tablets each day.

Because Entacapone Viatris increases the action of levodopa, it can also increase some of its side effects, especially at the start of treatment. To prevent this from happening, your doctor may lower your usual dose of levodopa for a few days or weeks. However, it is still important that you take an Entacapone Viatris tablet each time you take a dose of levodopa.

### ***How to take it***

**Swallow the tablet whole with a full glass of water at the same time as you take a dose of levodopa.**

If your stomach is upset after taking the tablet, take it with a meal or after a snack.

It does not matter if you take this medicine with or without food.

**Take your medicine at about the same times each day.**

Taking it at the same times each day will have the best effect. It will also help you remember when to take it.

**If you take iron supplements, take the iron supplement at least 2 to 3 hours before or after your dose of Entacapone Viatris.**

### ***How long to take it***

**Do not stop taking your medicine or change the dosage without checking with your doctor.**

This medicine helps to control your condition, but does not cure it. You may need to take Entacapone Viatris for a long time to control your symptoms.

If your doctor wants you to stop taking Entacapone Viatris, he or she may want you to gradually reduce the amount of Entacapone Viatris you take before stopping the medicine completely.

If you stop taking Entacapone Viatris, your doctor may increase the dose of your other medicines for Parkinson's disease to prevent your symptoms from getting worse.

### ***If you forget to take it***

**If you forget to take a tablet of Entacapone Viatris with your dose of levodopa, wait until the next dose of levodopa is due and start taking Entacapone Viatris again at this time.**

**Do not take a double dose to make up for the dose that you missed.**

This may increase the chance of you getting an unwanted side effect.

**If you are not sure what to do, ask your doctor or pharmacist.**

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

---

## **While you are taking Entacapone Viatris**

---

### ***Things you must do***

**If you are to be started on any new medicine, remind your doctor and pharmacist that you are taking Entacapone Viatris.**

**Tell any other doctors, dentists, and pharmacists who treat you that you are taking this medicine.**

**If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine.**

It may affect other medicines used during surgery.

**If you become pregnant while taking this medicine, tell your doctor immediately.**

**If you are about to have any blood tests, tell your doctor that you are taking this medicine.**

It may interfere with the results of some tests.

**Keep all of your doctor's appointments so that your progress can be checked.**

Your doctor may do some tests from time to time to make sure the medicine is working and to prevent unwanted side effects.

## ***Things you must not do***

**Do not take Entacapone Viatris to treat any other complaints unless your doctor tells you to.**

**Do not give your medicine to anyone else, even if they have the same condition as you.**

**Do not stop taking Entacapone Viatris suddenly, or change the dosage, without first checking with your doctor.**

Stopping Entacapone Viatris suddenly can cause some people to have side effects.

## ***Things to be careful of***

**Be careful driving, operating machinery or doing jobs that require you to be alert until you know how Entacapone Viatris affects you.**

This medicine may cause dizziness, light-headedness, sleepiness or drowsiness in some people. If you have any of these symptoms, do not drive, operate machinery or do anything else that could be dangerous.

**If you feel light-headed, dizzy or faint when getting out of bed or standing up, get up slowly.**

These symptoms may be a sign of low blood pressure. Standing up slowly, especially when you get up from bed or chairs, will help your body get used to the change in position and blood pressure. If this

problem continues or gets worse, talk to your doctor.

---

## **In case of overdose**

---

### ***If you take too much (overdose)***

**Immediately telephone your doctor or the National Poisons Information Centre (0800 POISON or 0800 764 766) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much Entacapone Viatris. Do this even if there are no signs of discomfort or poisoning.** You may need urgent medical attention.

Symptoms of an overdose may include sleepiness, confusion, decreased activity, low muscle strength and itchy rash.

---

## **Side effects**

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Entacapone Viatris.**

Like all medicines, Entacapone Viatris may have unwanted side effects in some people. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

**Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor or pharmacist if you notice any of the following and they worry you:**

- nausea (feeling sick) or vomiting
- diarrhoea or constipation
- aches and pains including abdominal pain
- dry mouth
- dizziness or light-headedness
- spinning sensation (vertigo)
- tiredness
- shakiness
- leg cramps
- headache
- trouble sleeping

- nightmares
- increased sweating
- a reddish-brown colour to the urine (this is known to happen with Entacapone Viatris and is usually harmless)
- discolouration of hair, beard, skin or nails.

The above list includes the more common side effects of your medicine.

**Tell your doctor as soon as possible if you notice any of the following:**

- unusual and uncontrolled movements of the body such as twisting, jerking or writhing movements
- extreme sleepiness or sudden onset of sleep in the middle of daytime activities
- worsening of your Parkinson's disease
- feeling depressed, confused or agitated
- hallucinations (seeing, hearing or feeling things that are not there)
- behavioural changes like the urge to gamble, compulsive spending/buying or increased sexual desire urges
- binge/compulsive eating
- signs of possible liver problems such as loss of appetite, feeling generally unwell, fever, itching
- signs of possible anaemia (low amount of iron in your blood) such as tiredness, headaches, shortness of breath when exercising, dizziness and looking pale
- progressive loss of appetite and asthenia (weakness, exhaustion) and weight decrease in a short period of time. Your doctor may decide to conduct a general medical evaluation including liver function.

The above list includes serious side effects that may require medical attention.

**If any of the following happen, tell your doctor immediately or**

**go to Accident and Emergency at your nearest hospital:**

- jaundice (yellowing of the skin or eyes)
- skin rash, itching or hives
- swelling of the face, lips, tongue or other parts of the body
- shortness of breath, wheezing or troubled breathing
- chest pain
- symptoms such as sudden increase in body temperature, sweating, fast heartbeat, muscle stiffness, extremely high blood pressure or severe convulsions. These may be symptoms of neuroleptic malignant syndrome (NMS)
- temporary paralysis or severe weakness of the muscles
- prolonged diarrhoea or diarrhoea with blood, mucus, stomach pain and fever.

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

**Tell your doctor or pharmacist if you notice anything that is making you feel unwell.**

Other side effects not listed above may also occur in some people.

Some of these side effects (for example, anaemia, changes in liver function and blood pressure) can only be found when your doctor does tests from time to time to check your progress.

**Do not be alarmed by the lists of possible side effects. You may not experience any of them.**

---

## After taking Entacapone Viatris

---

### Storage

**Keep your tablets in the bottle until it is time to take them.**

If you take the tablets out of the bottle they may not keep well.

**Keep your tablets in a cool dry place where the temperature stays below 25°C.**

Do not store Entacapone Viatris or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car. Heat and dampness can destroy some medicines.

**Keep it where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### Disposal

**If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.**

---

## Product description

---

### What it looks like

Entacapone Viatris tablets are light orange coloured, oval shaped, biconvex, film-coated tablets.

They are debossed with “EE200” on one side of the tablet and “M” on the other side.

Entacapone Viatris tablets are supplied in bottles of 100 tablets.

### Ingredients

Active ingredient:

Entacapone Viatris contains 200 mg of entacapone as the active ingredient.

Inactive ingredients:

Entacapone Viatris also contains in the tablet core:

- microcrystalline cellulose
- mannitol
- low-substituted hydroxypropylcellulose
- magnesium stearate
- hydrogenated vegetable oil.

The film-coating on the Entacapone Viatris tablet also contains:

- HPMC 2910 / hypromellose 6cP
- titanium dioxide
- glycerin
- magnesium stearate
- iron oxide yellow
- sucrose
- polysorbate 80

- iron oxide red.

Contains sulfites and gluten from wheat.

---

## If you want to know more

---

Should you have any questions regarding this product, please contact your pharmacist or doctor.

### Who supplies this medicine

Entacapone Viatris is supplied in New Zealand by:

Viatris Ltd  
PO Box 11183  
Ellerslie  
AUCKLAND  
[www.viatris.co.nz](http://www.viatris.co.nz)  
Telephone: 0800 168 169

---

## Date of Preparation

---

01 August 2024  
(Based on datasheet dated 01 August 2024)